

Natural Family Planning (NFP) Methods vs. Contraception

Methods	Possible Side Effects	% Effectiveness Avoiding Pregnancy	Harmful to Future Fertility	Responsible for Use
Natural Family Planning	Only method of FP that can be used to achieve/avoid pregnancy	Perfect Use †		Fosters Communication
Billings Ovulation Method - mucus based	None	98	No	Man and Woman
Creighton Model Fertility Care System - mucus based	None	99.5	No	Man and Woman
Couple to Couple League (CCL) - symptothermal, mucus & temperature based	None	99.4	No	Man and Woman
Marquette Model - mucus & electronic fertility monitor based	None	98	No	Man and Woman
SymptoPro Fertility Education - symptothermal, mucus & temperature based	None	99	No	Man and Woman
Barrier Methods*				
Condom	Allergic reaction or irritation.	98	No	Man
Diaphragm	Allergic reaction or irritation, Sometimes difficult to remove.	94	No	Woman
Vaginal Spermicides*				
Creams, Gels, Suppositories	Allergic reaction or irritation.	82	No	Woman
Sterilization Surgeries*				
Tubal Ligation	Surgery may be complicated by bleeding and/or infection. Increased rate: hysterectomy, ectopic pregnancy, painful periods.	99.5	Yes	Woman
Vasectomy	Surgery may be complicated by bleeding and/or infection. May increase risk of prostate cancer.	99.9	Yes	Man
Intrauterine Device (IUD) * **				
Affects sperm motility/survival and affects the lining of the uterus and prevents the implantation of the early human life in the uterus; <i>requires replacement every 10 years.</i>	Heavy, irregular or painful menses; painful intercourse; infection; ectopic pregnancy; perforation of the uterus.	99.4	Possible	Woman
Hormonal Contraception (2 hormones) - Pill, Patch, Vaginal Ring				
*Contraceptive: Suppresses ovulation (the release of an egg). If ovulation occurs, the sperm may be prevented from reaching the egg because of thickened cervical mucus. **Abortifacient: if ovulation occurs (2-29% of cycles) and conception occurs, the hormones act by preventing the implantation of the early human life in the uterus.	Common: breast tenderness/cysts, depression, weight gain, headaches, irregular bleeding. Rare: increased risk of breast cancer, liver tumors, stroke, heart attack, blood clots, elevated blood pressure.	99.6	May Delay	Woman
Hormonal Contraception (1 hormone = Progestin only)				
*Contraceptive and **Abortifacient: works like the combination hormonal contraception, but breakthrough ovulation occurs more frequently				
Implanon (Implant in skin) <i>replacement required every 3 years</i> ; breakthrough ovulation can occur after 30 months (3% between 30-36 months)	Irregular bleeding; acne; weight gain; headaches; depression, vaginal dryness.	99.9	May Delay	Woman
Mirena (IUD with hormone) <i>replacement required every 5 years</i> ; breakthrough ovulation 45% at 1 year and 75% at 4 years		99.8	May Delay	Woman
Progestin only Pill (breakthrough ovulation 33-65%)		99.6	May Delay	Woman
Depo-Provera (<i>injection every three months</i>) breakthrough ovulation can occur but incidence not known	See above; delays return of normal ovarian function by average of 8 months after last shot; decrease in bone density	99.7	Delays	Woman

Conception: union of sperm and egg; the beginning of human life

***Contraceptive:** prevents conception

****Abortifacient:** agent that ends human life after conception has occurred

† **Perfect Use:** method taught and practiced correctly

Credit:

Peter Danis, Family Physician, Certified FertilityCare Medical Consultant,
Gavin Puthoff, Ob/Gyn, Certified FertilityCare Medical Consultant, 2019

Natural Family Planning (NFP) Methods vs. Contraception 2019 REFERENCES

1. Hilgers TW, Stanford JB. Creighton Model NaproEducation Technology for Avoiding Pregnancy. *Journal of Reproductive Medicine* 1998; 43:495-502.
2. World Health Organization. A Prospective Multicenter Trial of the Ovulation Method of Natural Family Planning. II. The Effectiveness Phase. *Fertility and Sterility* 1981; 36 (5): 591-98.
3. Frank-Herrmann P, Heil J, Gnoth C, Toledo E, Baur S, Pyper C, Jenetzky E, Strowitzki T, Freundl G. The effectiveness of a fertility awareness based method to avoid pregnancy in relation to a couple's sexual behavior during the fertile time: a prospective longitudinal study. *Human Reproduction* 2007; 22: 1310–1319.
4. Fehring R, Schneider M, Raviele K. Efficacy of hormonal fertility monitoring as a method of natural family planning. *J Obstet Gynecol Neonat Nurs* 2007; 36:152–60.
5. Manhart MD, Duane M, Lind A, Sinai I, Golden-Tevald JDO. Fertility awareness-based methods of family planning: A review of effectiveness for avoiding pregnancy using SORT. *Osteopathic Family Physician* (2013) 5, 2–8.
6. Rulin MC, Davidson AR, Philliber SG. Long term effect of tubal sterilizations on menstrual indices and pelvic pain. *Obstetrics and gynecology* 1993; 82: 118-21.
7. Peterson HB, Zhisen X, Hughes JM, et al. The risk of ectopic pregnancy after tubal sterilization. *The New England Journal of Medicine* 1997; 336: 72-7.
8. Siddiqui MM, Wilson KM, Epstein MM, Rider JR, Martin NE, Stampfer MJ, Giovannucci EL, Mucci LA. Vasectomy and Risk of Aggressive Prostate Cancer:A 24-Year Follow-Up Study. *J Clin Oncol* 2014; 32:3033-3038.
9. Larimore WL, Stanford JB. Postfertilization Effects of Oral Contraceptives and Their Relationship to Informed Consent. *Archives of Family Medicine* 2000; 9: 126-133.
10. Hassan MA, Killick SR. Is previous use of hormonal contraception associated with a detrimental effect on subsequent fecundity? *Human Reproduction* 2004; 19: 344-351.
11. Kahlenborn C, Modugno F, Potter DM, Severs WB. Oral Contraceptive Use as a Risk Factor for Premenopausal Breast Cancer: A Meta-analysis. *Mayo Clin Proc* 2006; 81:1290-1302.
12. Mørch LS, Skovlund CW, Hannaford PC, et al. Contemporary Hormonal Contraception and the Risk of Breast Cancer. *NEJM* 2017;377: 2228-39.
13. Skovlund CW, Steinrud Mørch LS, Kessing LV, Lidegaard O. Association of Hormonal Contraception with Depression. *JAMA Psychiatry*. 2016;73(11):1154-1162
14. Etmnan M, Takkouche B, Isorna FC, Samii A. Risk of ischaemic stroke in people with migraine: systematic review and meta-analysis of observational studies. *BMJ* 2004, doi:10.1136/bmj.38302.504063.8F (published 20 December 2004)
15. Hatcher RA, et al. Contraceptive Technology. *Ardent Media Inc.*, 2018.